



## Guaranteed Minimum Analysis

Salicylic Acid.....	72.5%
Potassium (available K <sub>2</sub> O).....	27.5%

**EZ-Gro-SA-P-72.5%** contains Salicylic Acid, a naturally occurring plant hormone, which plays a vital role in plant growth and development, photosynthesis, transpiration, ion uptake and transport. Salicylic Acid improves resistance to environmental stressors including drought, temperature, and salinity; and enhances flowering and fruit yield.

## APPLICATION INSTRUCTIONS:

**EZ-Gro-SA-P-72.5%** can be used as a foliar spray, a soil drench and through fertigation systems. | It can be mixed with most fungicides, insecticides and other liquid fertilizers. Do not mix concentrates together. Add water to spray tank to ¾ full followed by the **EZ-Gro-SA-P-72.5%**, pesticides and fertilizers. | When mixing with dry or liquid fertilizer, do not exceed the recommended rate of 7 g/ha/application.

**To make a 10% concentrated solution: dissolve 140 grams into 1 litre of water (20 oz/1 gallon).**

## DANGER

### Keep out of reach of children and animals:

Harmful if swallowed. | Causes serious eye damage.  
| Avoid breathing dust, spray or mist. Use only outdoors or in a well-ventilated location. | Wear protective gloves, clothing, eye and NIOSH/OSHA approved respirator during product mixing and loading. | Do not eat, drink or smoke when using this product. | Wash thoroughly after handling.  
| Avoid entry into water sources. Do not let product enter drains when rinsing or disposing of equipment wash water.  
| Immediately contact a health care provider if exposed, concerned or feeling unwell. | If swallowed, rinse mouth.  
| If in eyes, rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. | If inhaled, move person to fresh air. | Store locked up in cool, dry, well-ventilated place out of direct sunlight. | Dispose of contents and packaging in accordance with local, regional and federal requirements.

# EZ-GRO-SA-P-72.5%

## APPLICATION CHART:

CROP	APPLICATION RATE		TIMING / FREQUENCY
	10% Solution	Undiluted	
Nursery Stock (Containerized and field grown): deciduous and Evergreen trees, annuals, perennials, herbs, and vegetables	40 ml / 100 L water (1.25 oz / 20 ga water)	N/A	Apply every 30 days or as needed commencing at transplant or 3-5 leaf stage.
Fruits and Berries: plums, apricots, cherry, peach, blueberries, blackberries, raspberries, strawberries			3-4 applications commencing at 75% petal drop and repeating with 21-28 day intervals
Apples and Pears			3-4 applications commencing at 50% flowering and repeating with 21 day intervals
Fruiting vegetables: tomatoes, peppers, beans, peas, onions	50 ml / ha (0.7 oz / acre)	7 g / ha (0.1 oz / acre)	First application prior to fruit set and repeat after 2 weeks
Lettuce, cauliflower, cabbage, Chinese cabbage			First application at 3-5 leaf stage and repeat after 2 weeks
Potatoes			First application at tuber initiation and repeat after 2 weeks
Cucurbits: pumpkins, squash, melons, cucumbers			First application at flowering and repeat after 21 days
Grapes: table and wine			Apply every 4-6 weeks commencing once plants begin to grow
Herbs: basil, chives, coriander, dill, parsley, sage			Apply every 4-6 weeks commencing once plants begin to grow
Field crops: barley, corn, oats, soybean, sugar beet, wheat			After plants reach the 3-5 leaf stage. Use with sufficient water to cover.
Grasses grown for seed, sod, pasture, forage, alfalfa			Apply when 2-4" tall. Use with sufficient water to cover.
Compost and Soil	40 ml / 100 L of soil/compost	N/A	Apply to soil or compost and mix thoroughly.
Hydroponics and Medical Marijuana	N/A	1 g/200 L water	Apply to roots weekly through grow and bloom, beginning at transplant

- ☐ 1 kg / 2.2 lb
- ☐ 5 kg / 11 lb
- ☐ 10 kg / 22 lb